

S.C. Zoltai Canadian Forestry ServiceC. Tarnocai Manitoba Soil SurveyE.T. Oswald Canadian Forestry Service

Edited and produced by


J. Thie Canada Land Inventory Project for Manitoba

CROMORANT LAKE BIOPHYSICAL PROJECT

CRANVERRY PORTAGE, MANITOBA JUNE 17-20/ 1970

TABLE OF CONTENTS

																																							P	age
INTRO	DUC	CTION			•			•								•					٠.		•																	3
STOP	1,	Plot	1	٠.																			•													•				5
		Plot	2	Α.	•		٠.	•	٠.												٠.	•	•								•			• •	•	•	 •			7
		Plot	2	В.	•					•	•		•			•		•		•											•			٠.		•	 •			9
		Plot	2	С.	•	• •		•	٠.		•		•	٠.		•		•		•		•	•		•				•		•		•			•	 •			10
		Plot	3	٠.	•			•	٠.	•	•		•			•		•		•		•	•	٠.	•		•	• •	•		•	٠.	•			•	 •			12
STOP	2,	Plot	4		•																																			14
STOP	3,	P1ot	5																				•														 •			18
STOP	4,	Plot	6	٠.																																	 •			22
STOP	5,	Plot	7																	•																				25
STOP	6,	P1ot	8																																					28
		Plot	9	١			•				•									•								٠.												30
STOP	7,	Plot	1	.0.																																	 •			33
STOP	8,	Plot	1	.1																																				36
STOP	9,	Plot	1	.2.																																		٠.		39
APPE	NDI	ХΙ-	M	lec	h	an	ii	ca	1	a	n	d	C	he	em	ii	ca	1	Α	۱n	a]	Ly	S	is	; (of		Sc	i	1s							 			41



The Cormorant Lake mapsheet lies north of the 54° parallel, at the same latitude as Hamilton Inlet in Labrador, Cape Henrietta Maria in Ontario and northern Queen Charlotte Islands in British Columbia. The climate is cold continental, characterized by relatively short, cool growing season and moderate precipitation. Permafrost may be found in some peatlands throughout the mapsheet.

Physiographically the area is divided into two almost equal portions. The southern half is vastly different from the northern half. In the south the land is quite flat, interrupted only by some low Ordovician dolomite plateaus and drumlins. The northern half is on the Precambrian Shield and is dominated by broken, locally rugged bedrock topography and numerous bedrock outcrops. Lakes are very numerous, especially in the north, where they tend to be relatively deep. Many of the lakes in the southern half are very shallow.

Vegetation throughout the area is composed of boreal forests, with black spruce as the dominant species. Jack pine forests are induced by fires on the drier sites. Trembling aspen-white spruce forests occur mainly in the southwestern part of the map area and along the contact of the two physiographic provinces. Bogs and fens cover extensive areas, especially in the south. Exploitation of the forests has been accelerated since 1968, in preparation for a forest industry complex just south of the map area.

Moose is the most abundant ungulate, but woodland caribou also occurs in the northern half of the area. Moose attract many hunters during the hunting season.

The large number of picturesque lakes lure many fishermen and tourists to the wilderness areas. Walleye, northern pike and Lake trout are the main

sports fish species. The Grass River and Clearwater Lake Provincial Parks provide camping facilities for families and fishing or hunting parties.

Mining is the economic mainstay of the area. Flin Flon and Snow Lake are important mining and refining centres. A large base metal deposit was recently discovered beneath Lake Athapapuskow.

The purpose of the field trip is to acquaint visitors with the area by pointing out the features which make this area unlike any other. During discussions this information can be drawn together within the Bio-Physical Land Classification System, and related to other parts of the country. Throughout the field trip the needs of users of this information will be emphasized.



Stereotriplet A. A13879:-99-101, fall 1953. Scale ±1/68,000. T.65, R.28W.

ATHAPAPUSKOW Landscape Unit (293 sq. mi.)

Stereotriplet A

Moderately to weakly broken area of shallow sandy till over volcanic bedrock in northern part, and deep to shallow loamy till over dolomitic bedrock in the southern part, with lacustrine clay in some depressions.

Lakes occupy 38% of the area. Lakes of various size classes cover the following proportions of the landscape unit:

Large lakes:	34%
Relatively small lakes:	1%
Small lakes:	3%

The lakes are moderately deep to deep with irregular to very irregular shorelines in the north, but only somewhat irregular shorelines in the south. The shore material is mainly bedrock or bouldery till, but broad sandy beaches do occur along the southern shore of Lake Athapapuskow. The lakes are drained by rivers or small creeks. Some small lakes in the south drain through bogs.

Land Unit CS-62b

Moderately broken area of shallow silty sand till over volcanic bedrock (55%) with bare volcanic bedrock exposures (25%). Minor areas of deep, low base silty sand in some valleys; deep fibric peat in some valleys; shallow to deep moderately calcareous lacustrine clay in some valleys.

PLOT #1 Land Upper slope on a crag-and-tail structure, consisting

Stereotriplet A+B of deep low base silty sand till. Moisture regime

is fresh (2), or well drained.

Soil Fay Lake Series

The Fay Lake series consists of well drained Degraded

Dystric Brunisol soils developed on non-calcareous

sandy loam to loamy sand till derived from Precambrian granitic and basaltic rocks.

- L-F 3 to 0 inches, very dark grey (10YR 3/1, moist), forest litter of needles and small branches; extremely acid; abrupt, smooth boundary.
- Ae 0 to 4 inches, greyish brown (10YR 5/2, moist), loamy sand; single grained; loose when moist, loose when dry; non-sticky, non-plastic; strongly acid; clear, wavy boundary.
- Bm 4 to 12 inches, yellowish brown (10YR 5/6, moist), loamy sand; single grained; loose when moist, loose when dry; non-sticky, non-plastic; medium acid; clear, wavy boundary.
- BC 12 to 24 inches, light brownish grey (10YR 6/2, moist), loamy sand; single grained; loose when moist, loose when dry; non-sticky, non-plastic; medium acid; clear, smooth boundary.
- C 24 to 29 inches, light grey (10YR 7/2, moist), loamy sand; single grained; loose when moist, loose when dry; non-sticky, non-plastic; slightly acid.


Analytical data in Table 1, Appendix I.

Forest Black spruce - jack pine, M.A.I. = 36 cu.ft./ac./yr.

	<u>bS</u>	<u>jP</u>
Diameter	8.2 in.	9.0 in.
Height	60 ft.	55 ft.
Age	115 yrs.	105 yrs.
Basal area	120 sq. ft.	60 sq. ft.

Vegetation

Trees - Pinus banksiana
Picea mariana
Betula papyrifera



STOP 1 Plots 1,2,3

Stereopair B. A20040:132-133, 1967. Scale $\pm 1/16,000$. T.65, R.28W. Area covered by this Stereopair indicated on Stereotriplet A.

Forbs - Aralia hispida

Epilobium angustifolium

Linnea borealis

Pyrola secunda

Rubus idaeus

Bryophytes & lichens

(Open, sandy)

Ceratodon purpureus

(Litter)

Oncophorus polycarpus Var. strumiferus

Pohlia nutans

Ptilidium ciliare

Capability Classification

Forestry: 5 F

Agriculture: 6º

Wildlife: 4F

Recreation: 600E

PLOT #2a <u>Land</u> Top of low base (volcanic) bedrock outcrop. Moisture Stereotriplet A+B regime: very dry (0) or rapidly drained.

Soil Rock Outcrop

The Rock Outcrop consists of 0 to 4 inches of rapidly drained surface deposits over Precambrian bedrock.

L-H - 1 to 0 inches, very dark greyish brown (10YR 3/2, moist), forest litter; strongly acid; abrupt, smooth boundary.

R - 0+ inches, basaltic bedrock.

Analytical data in Table 2, Appendix I.

Forest Open, scattered black spruce - jack pine.

Vegetation

Trees - Pinus banksiana

Picea mariana

Betula papyrifera

Tall shrubs - Alnus crispa

Juniperus communis

Low shrubs - Arctostaphylos uva-ursi

Vaccinium angustifolium

Forbs - Anemone patens

Fragaria virginiana

Heracleum lanatum

Potentilla tridentata

Saxifraga tricuspidata

Bryophytes & lichens - Cladonia spp.

(Litter) Pohlia, cf. bulbifera

P. nutans

Ceratodon purpureus

Drepanocladus unicinatus

(Rock crevices) Oncophorus polycarpus var. strumiferus

O. virens

Mnium spinulosum

Ptilidium ciliare

(Under rock ledges) Distichum capillaceum

Pohlia sp.

(On rocks) Cratoneuron falcatum

Oncophorus polycarpus var. strumiferus

Pohlia nutans

Drepanocladus fluitans

Mnium spinulosum

Ptilidium ciliare

Lophozia ventricosa

Capability Classification

Forestry: 7 %

Agriculture: 7^R

Wildlife: 3° 6 k

Recreation: 60E0

PLOT #2b Land Depression in bedrock filled by shallow low base Stereotriplet A+B

till. Moisture regime is very moist (5), or

imperfectly drained.

Athapap Series Soi1

> The Athapap series consists of imperfectly drained Gleyed Lithic Dystric Brunisol soils developed on non-calcareous sandy loam to loamy sand till derived from Precambrian granitic and basaltic rocks, with a lithic contact between 4 to 20 inches of the mineral surface.

L-H - $1\frac{1}{2}$ to 0 inches, very dark grey (10YR 3/1, moist), forest litter of needles and small branches; extremely acid; abrupt, smooth boundary.

- 0 to 4 inches, dark brown (10YR 4/3, moist), sandy Bmloam; single grained; loose when moist, loose when dry; non-sticky, non-plastic; very strongly acid; clear, smooth boundary.

- 6 to 14 inches, olive (5Y 5/3, moist), sandy loam; Cg single grained; loose when moist, loose when dry; non-sticky, non-plastic; strongly acid; abrupt, smooth boundary.

- 14+ inches, basaltic bedrock. R

Analytical data in Table 3, Appendix I.

Black spruce, M.A.I. = 24 cu.ft./ac./yr. Forest

> bS Diameter 4.3 in. Height 48 ft. Age 115 yrs. Basal area 140 sq. ft.

Vegetation

Trees - Picea mariana
Pinus banksiana

Betula papyrifera

Tall shrubs - Alnus sp.

Rosa sp.

Low shrubs - Ledum groenlandicum

Vaccinium vitis-idaea

Forbs - Linnaea borealis

Pyrola asarifolia

P. secunda

Trientalis borealis

Lichens - Cladonia sp.

Capability Classification

Forestry: 6w

Agriculture: 6 %

Wildlife:

6 4°

Recreation: 60EQ

PLOT #2c Land Low plateau, shallow stony lacustrine clay over

Stereotriplet A+B bedrock. Moisture regime: moist (4), or imperfectly drained.

Soil Herb Lake Series

The Herb Lake series consists of imperfectly drained Gleyed Lithic Grey Luvisol soils developed on strongly calcareous lacustrine clay, with a lithic contact between 4 to 20 inches of the mineral surface.

L-H - 3 to 0 inches, very dark brown (10YR 2/2, moist), forest litter of needles and small branches; extremely acid; abrupt, smooth boundary.

- Ae 0 to 3 inches, grey (10YR 5/1, moist), clay loam; moderate, medium platy; friable when moist, hard when dry; slightly sticky and plastic; very strongly acid; clear, smooth boundary.
- ABg 3 to 5 inches, greyish brown (10YR 5/2, moist), clay; weak, medium, coarse columnar, breaking to strong, coarse subangular blocky; firm when moist, very hard when dry; sticky and plastic; strongly acid; clear, wavy boundary.
- Btg 5 to 10 inches, dark brown (10YR 4/3, moist), clay; moderate, medium subangular blocky; firm when moist, very hard when dry; sticky and plastic; strongly acid; abrupt, smooth boundary.

R - 10+ inches, basaltic bedrock.

Analytical data in Table 4, Appendix I.

Forest Black spruce, M.A.I. = 26 cu.ft./ac./yr.

bS
Diameter 4.4 in.
Height 44 ft.
Age 119 yrs.
Basal area 150 sq. ft.

Vegetation

Trees - Picea mariana
Pinus banksiana
Betula papyrifera

Tall shrubs - Alnus sp.

Rosa sp.

Low shrubs - Ledum groenlandicum

Vaccinium vitis-idaea

Forbs - Linnaea borealis

Pyrola asarifolia

P. secunda

Trientalis borealis

Trientalis borealis

11

Lichens - Cladonia sp.

Capability Classification

Forestry: 5w

Agriculture: 5 R

Wildlife: 5 R 4G

Recreation: 60E0

PLOT #3
Stereotriplet A+B

<u>Land</u> Peat plateau: perennially frozen fibric to mesic peat, with collapse scars. Moisture regime: moist (4), or imperfectly drained.

Soil Batty Lake Series

The Batty Lake series consists of imperfectly drained Cryic Mesic Fibrisol soils developed on forest peat.

Of - 0 to 37 inches, very dark grey (10YR 3/1, moist), slightly woody, fibrous feathermoss peat; very strongly acid; fiber content approximately 73 percent.

Omz - 37 to 170 inches, black (10YR 2/1, moist), frozen, moderately decomposed, mixed feathermoss and woody peat with segregated ice crystals and ice lenses; strongly acid; fiber content approximately 57 percent.

Om - 170+ inches, black (10YR 2/1, moist), moderately well decomposed, mixed feathermoss and woody peat.

Analytical data in Table 5, Appendix I.

Forest Black spruce, M.A.I. = 9 cu.ft./ac./yr.

 $\frac{bS}{a}$ Diameter 3.7 in.

Height 25 ft.

Age 118 yrs.

Basal area 100 sq. ft.

Vegetation

Trees - Picea mariana

Low shrubs - Ledum groenlandicum

Vaccinium vitis-idaea

Forbs - Drosera rotundifolia

Maianthemum canadense

Oxycoccus microcarpus

Rubus chamaemorus

Bryophytes & lichens - Cladonia spp.

(On litter) Pleurozium schreberi

Sphagnum fuscum

Sphagnum warnstorfianum

Tetraplodon angustatus

Meesia uliginosa

Capability Classification

Forestry: 7 %

Agriculture: O

Wildlife:

6 5 3°

Recreation: 60E0

SIMONHOUSE Landscape Unit (124 sq. mi.)

Stereotriplet C

Weakly to very weakly broken area of deep loamy till, with some plateaus of shallow loamy till over bedrock, and lacustrine clay till in some valleys. In the northern, Canadian Shield area shallow sandy till over bedrock and bare bedrock ridges dominate.

Lakes cover 44% of the area. Lakes of various size classes occupy the following proportion of the landscape unit:

Large lakes:	41%
Relatively small lakes:	2%
Small lakes:	1%

The larger lakes have irregular shorelines with many islands in the Precambrian portion of their basin. These lakes are moderately deep and are drained by small creeks or through straits. The smaller lakes in the south have regular shorelines and are shallow, often without open outlets. The shore material is generally bouldery till or, locally, organic matter. Broad sandy beaches occur on Simonhouse Lake.

Land Unit ML-72L


Weakly broken area of slopes and flats of deep somewhat calcareous sandy loam (70%) with plateaus of shallow highly calcareous loam over dolomitic bedrock (20%). Minor areas of deep to shallow mesic or fibric peat in some depressions.

PLOT #4 Land Middle slope on a low drumlin, weakly calcareous

Stereotriplet C
Stereopair E

loamy till. Moisture regime: fresh (2), or well

drained.



Stereotriplet C. A13904:67-69, fall 1953. Scale ±1/68,000. T.64, R.26. For detail see Stereopairs D and E.

Soil Egg Lake Series

The Egg Lake series consists of well drained Orthic Grey Luvisol soils developed on moderately to very strongly calcareous, sandy loam to loamy till.

- L-H 1½ to 0 inches, very dark grey (10YR 3/1, dry), black (10YR 2/1, moist), forest litter; medium acid; abrupt, smooth boundary.
- Ae 0 to 8 inches, light grey (2.5Y 7/2, dry), dark greyish brown (10YR 4/2, moist), sandy loam; moderate, fine platy; friable when moist, slightly hard when dry; non-sticky, non-plastic; medium acid; abrupt, smooth boundary.
- Bt 8 to 14 inches, yellowish brown (10YR 5/4, dry), dark brown (7.5YR 3/2, moist), sandy loam; moderate, medium subangular blocky; firm when moist, very hard when dry; slightly sticky and slightly plastic; mildly alkaline; clear, smooth boundary.
- BC 14 to 19 inches, light grey (10YR 7/1, dry), yellowish brown (10YR 5/4, moist) sandy loam; moderate, medium subangular blocky; firm when moist, very hard when dry; slightly sticky and slightly plastic; mildly alkaline; moderately calcareous; gradual, wavy boundary.
- Cca 19 to 34 inches, light grey (10YR 7/1, dry), light brownish grey (10YR 6/2, moist), sandy loam; moderate, fine subangular blocky; firm when moist, very hard when dry; slightly sticky and slightly plastic; mildly alkaline; strongly calcareous; clear, smooth boundary.
- C 34+ inches, light grey (10YR 7/1, dry), light brownish grey (10YR 6/2, moist), sandy loam; moderate, fine subangular blocky; firm when moist, very hard when dry; mildly alkaline; moderately calcareous.

Forest Aspen-white spruce, M.A.I. = 53 cu.ft./ac./yr.

	<u>tA</u>	<u>wS</u>
Diameter	6.7 in.	6.4 in.
Height	48 ft.	32 ft.
Age	55 yrs.	48 yrs.
Basal area	120 sq. ft.	50 sq. ft.

Vegetation

Trees - Picea glauca
Picea mariana
Populus tremuloides

Tall shrubs - Alnus crispa Rosa spp.

Low shrubs - Arctostaphylos uva-ursi
Symphoricarpos sp.
Vaccinium vitis-idaea
Viburnum edule

Forbs - Aralia nudicaulis

Epilobium angustifolium

Fragaria virginiana

Linnaea borealis

Maianthemum canadense

Pyrola secunda

Rubus pubescens

Bryophytes & lichens

(On granitic rock) Drepanocladus uncinatus
Plagiothecium turfaceum
Brachythecium flexicaule
Mnium cuspidatum

(On base of aspen) Pylaisia polyantha

Brachythecium salebrosum

(On rotten stump) Pohlia bulbifera

Capability Classification

Forestry: 4

Agriculture: 5^F

Wildlife: 2° Recreation: 5°

GOOSE Landscape Unit (196 sq. mi.)

Stereotriplet C

Weakly to very weakly broken area of deep loamy till with some shallow till over bedrock, and deep lacustrine clay and clay till in low-lying areas. Peat occurs in most depressions.

Lakes cover 34% of the area. Lakes of various size classes occupy the following proportion of the landscape unit:

Large lakes: 26%
Relatively small lakes: 6%
Small lakes: 2%

The lakes are moderately deep, with large shallow bays. The shorelines are relatively regular, with only a few islands. The shore material is mainly bouldery till or bedrock, but locally organic matter may form the shores. The larger lakes all have outlets, but the small lakes are without open drainage.

Land Unit ML-P50

Weakly broken area of deep mesic peat (60%), with low knolls and flats of deep highly calcareous clay (15%) and loam (15%). Minor areas of shallow fibric peat over clay or loam in some depressions.

PLOT #5 Land Palsa: perennially frozen fibric to mesic peat,

Stereotriplet C
Stereopair D with unfrozen fens in depressions. Moisture regime:

moist (4), or imperfectly drained.

Soil Batty Lake Series

The Batty Lake series consists of imperfectly drained Cryic Mesic Fibrisol soils developed on forest peat.

- Of1 0 to 7 inches, very dark greyish brown (10YR 3/2, moist), slightly woody, fibrous feathermoss peat; layer of charcoal and ash; slightly acid; fiber content approximately 82 percent.
- Of2 7 to 34 inches, very dark grey (10YR 3/1, moist), slightly woody, fibrous feathermoss peat with thin layer of Sphagnum peat; neutral; fiber content approximately 81 percent.
- Omz 34 to 96 inches, black (10YR 2/1, moist), frozen, moderately decomposed, mixed forest and fen peat; alternating layers of marl in the lower portion of this layer; neutral; fiber content approximately 42 percent.
- IICgz 96 to 163 inches, greenish grey (5GY 5/1, moist), frozen clay; ice lenses 0.5 to 4 inches thick.
 - IICg 163 to 167 inches, greenish grey (5GY 5/1, moist),
 unfrozen, varved clay.

Analytical data in Table 7, Appendix I.

Forest Black spruce, M.A.I. = 9 cu.ft./ac./yr.

	bS	
Diameter	3.2	in.
Height	23	ft.
Age	83	yrs.
Basal area	70	sq. ft.

Vegetation

Trees - Picea mariana

Betula papyrifera

Tall shrubs - Betula glandulosa
Salix bebbiana

Low shrubs - Andromeda polifolia
Chamaedaphne calyculata
Ledum groenlandicum

Vaccinium vitis-idaea Oxycoccos microcarpus Empetrum nigrum Rubus chamaemorus

Forbs - Drosera rotundifolia Equisetum spp.

Bryophytes & lichens

(On litter)

Bryum inclinatum

Ceratodon purpureus

Dicranum bergeri

Bryoerythrophyllum recurvirostrum

(On tops of hummocks)

Ptilium crista-castrensis
Dicranum rugosum
Pleurozium schreberi
Cladonia rangiferina
C. mitis
C. alpestris

Dicranum bonjeani Sphagnum fuscum

(On sides of hummocks)

Sphagnum fuscum
Pleurozium schreberi
Aulacomnium plaustre
Tomenthypnum nitens
Thuidium microphyllum
Pohlia nutans

(In standing water)

Sphagnum fallax
S. rubellum
S. wulfianum
Calliergon giganteum
Aulacomnium palustre

Capability Classification

Forestry: 7_w^k

Agriculture: O

Recreation: $6^{\frac{M}{5}} 2^{\frac{G}{5}}$

WANLESS Landscape Unit (93 sq. mi.)

Stereotriplet F

Weakly to very weakly broken area of lacustrine clay, with low hills of deep loamy and clay till, and peat plains in depressions. A lake modified till escarpment occurs on the eastern margin.

Lakes occupy 2% of the area. Lakes of various size classes cover the following proportion of the landscape unit:

Relatively small lakes: 1%

Small lakes: 1%

The lakes are shallow and their shorelines are regular. The shores are formed by organic matter or by loam, in which case they are stony or bouldery. None of the lakes has an open outlet.

Land Unit ML-49h

Weakly broken area of deep highly calcareous clay flats and slopes (95%). Minor areas of deep to shallow mesic and fibric peat in depressions.

PLOT #6 <u>Land</u> Level highly calcareous lacustrine clay. Moisture Stereotriplet F regime: moist (3), or well drained.

Soil Wabowden Series

The Wabowden series consists of well drained Solodic Grey Luvisol soils developed on moderately to strongly calcareous lacustrine clay.

L-H - 2 to 0 inches, very dark grey (10YR 3/1, moist), forest litter; very strongly acid; abrupt, smooth boundary.



Stereotriplet F. A13870:57-59, fall 1953. Scale ±1/68,000. T.60,61, R.26W. For detail see Stereopair G.

- Ae 0 to 4½ inches, light grey (10YR 6/1, moist), clay; moderately coarse platy; friable when moist, hard when dry; slightly sticky and plastic; very strongly acid; clear, smooth boundary.
- AB 4½ to 9½ inches, light brownish grey (10YR 6/2, moist), clay; strong, coarse columnar; firm when moist, very hard when dry; sticky and plastic; strongly acid; gradual, wavy boundary.
- Bt 9½ to 18½ inches, greyish brown (2.5Y 5/2, moist), clay; strong, medium subangular blocky; firm when moist, hard when dry; sticky and plastic; slightly acid; gradual, wavy boundary.
- BC 18½ to 22½ inches, dark grey (10YR 4/1, moist), clay; strong medium subangular blocky; firm when moist, hard when dry; sticky and plastic; neutral; gradual, wavy boundary.
- C1 22½ to 30 inches, dark grey (10YR 4/1, moist), clay; moderate, medium subangular blocky; firm when moist, hard when dry; sticky and plastic; mildly alkaline; gradual, wavy boundary.
- C2 30 to 39 inches, grey (10YR 5/1, moist), clay; moderate, fine to medium subangular blocky; firm when moist, hard when dry; sticky and plastic; mildly alkaline.

Analytical data in Table 8, Appendix I.

Forest Jack pine-black spruce, M.A.I. = 62 cu.ft./ac./yr.

	jР	<u>bS</u>
Diameter	8.4 in.	8.5 in.
Height	65 ft.	65 ft.
Age	76 yrs.	70 yrs.
Basal area	100 sq. ft.	80 sq. ft.

Vegetation

Trees - Pinus banksiana

Picea mariana

Populus tremuloides

Tall shrubs

- Rosa sp.

Low shrubs

- Vaccinium angustifolium

Viburnum edule

Forbs

- Aralia nudicaulis

Cornus canadensis

Epilobium angustifolium

Linnea borealis

Maianthemum canadense

Mitella nuda Rubus idaeus

Bryophytes & lichens

(On old birch

stump)

Bryum pendulum

Brachythecium oxycladon

(On litter)

Pleurozium schreberi

Ptilium crista-castrensis

Capability Classification

Forestry: 4

Agriculture: 4º

Wildlife: 2°

Recreation: 50E



Stereotriplet H. A13870:115-112, fall 1953. Scale ±1/68,000. T.58,59, R.26W.

CLEARWATER Landscape Unit (232 sq. mi.)

Stereotriplet H

Weakly broken area of deep loamy till, with plateaus of shallow loamy till over bedrock and bare bedrock. Peat plains and lacustrine clay occupy many depressions.

Lakes occupy 45% of the area. Lakes of various size classes cover the following proportion of the landscape unit:

Large lakes: 43%
Small lakes: 2%

Clearwater Lake is moderately deep with some shallow bays, having a somewhat irregular shoreline. The smaller lakes have regular shorelines and are shallow. The shores are mainly bouldery or rocky, but some sandy beaches occur between headlands. Organic matter forms the shores of some small lakes. All but some small muskeg lakes have open outlets.

Land Unit ML-P45

Weakly broken area of shallow fibric peat over loam (60%) in depressions and lower slopes, with deep very highly calcareous loam (30%) on low hills. Minor areas of deep fibric peat in depressions; very highly calcareous gravelly sand in some low ridges.

PLOT #7 Land Lower slope of gently sloping drumlinoid structure,

Stereotriplet H shallow fibric peat over very highly calcareous loamy
till. Moisture regime: wet (8), or poorly drained.

Soil Dering Series Peaty Phase

The Dering series peaty phase consists of poorly
drained Rego Gleysol soils developed on extremely
calcareous loamy till.

L-F - 8½ to 4½ inches, brown (10YR 5/3, moist), fibrous sphagnum peat; extremely acid; abrupt, smooth boundary.

H - 4½ to 0 inches, very dark grey (10YR 3/1, moist), moderately well decomposed forest peat; neutral; abrupt, smooth boundary.

Ahg - 0 to 4 inches, black (10YR 2/1, moist), fine sandy loam; weak, fine subangular blocky; friable when moist, slightly hard when dry; sticky and slightly plastic; neutral; clear, smooth boundary.

Cg - 4 to 11 inches, light brownish grey (2.5Y 6/2, moist), silt loam; weak, fine subangular blocky; friable when moist, slightly hard when dry; sticky and slightly plastic; mildly alkaline.

Analytical data in Table 9, Appendix I.

Forest Black spruce, M.A.I. = 5 cu.ft./ac./yr.

Diameter 4.3 in.
Height 25 ft.
Age 195 yrs.
Basal area 80 sq. ft.

Vegetation

Trees - Picea mariana

Tall shrubs - Betula glandulosa
Salix bebbiana

Low shrubs - Andromeda polifolia
Arctostaphylos uva-ursi
Chamaedaphne calyculata
Juniperus horizontalis
Ledum groenlandicum
Vaccinium angustifolium
Vaccinium vitis-idaea
Oxycoccos microcarpus

Forbs - Arctostaphylos rubra
Drosera rotundifolia
Equisetum sp.

Bryophytes & lichens

(On litter) Pleurozium schreberi

Dicranum dummondii Sphagnum nemoreum


Capability Classification

Forestry: 7 w

Agriculture: 0

Wildlife: 6 3

Recreation: 60E



Cross section of Palsa along Line 4, showing permafrost core and the thickness of active layer related to vegetation cover. (From: Properties of a Wooded Palsa in Northern Manitoba by S.C. Zoltai and C. Tarnocai, in manuscript.

SIMONHOUSE Landscape Unit (124 sq. mi.)

Stereopair M

Weakly to very weakly broken area of deep loamy till, with some plateaus of shallow loamy till over bedrock, and lacustrine clay till in some valleys. In the northern, Canadian Shield area shallow sandy till over bedrock and bare bedrock ridges dominate.

Lakes cover 44% of the area. Lakes of various size classes occupy the following proportion of the landscape unit:

Large lakes	5		41%
Relatively	small	lakes:	2%

The larger lakes have irregular shorelines with many islands in the Precambrian portion of their basin. These lakes are moderately deep and are drained by small creeks or through straits. The smaller lakes in the south have regular shorelines and are shallow, often without open outlets. The shore material is generally bouldery till or, locally, organic matter. Broad sandy beaches occur on Simonhouse Lake.

Land Unit ML-36h

Weakly broken area of deep highly calcareous loam (60%) on slopes and flats, shallow loam over dolomitic bedrock (10%), with deep highly calcareous clay in depressions (10%). Minor areas of highly calcareous gravelly sand in some low ridges; shallow fibric peat over loam or clay in some depressions; deep mesic peat in some depressions.

PLOT #8 Land Gently sloping toe of beach ridge, moderately


Stereotriplet L
Stereopair M calcareous medium sand. Moisture regime: somewhat fresh (1), or well drained.

STOP 6 Plots 8,9



Stereopair M. A13789:202-203, 1953. Scale ±1/68,000. T.63, R.25W.

Stereotriplet L. A20040:209-211, 1967. Scale ±1/16,000. T.63, R.25. Detail of Sp.M.



Soil Clarke Series

The Clarke series consists of well drained Degraded Dystric Brunisol soils developed on glacio-fluvial sand deposits.

- L-H 2 to 0 inches, black (10YR 2/1, moist), forest litter of jack pine needles; very strongly acid; abrupt, smooth boundary.
- Ae 0 to 6 inches, light grey (10YR 7/1, moist), sand; single grained; loose when moist, loose when dry; non-sticky, non-plastic; strongly acid; abrupt, smooth boundary.
- Bm 6 to 14 inches, yellowish brown (10YR 5/6, moist), sand; single grained; loose when moist, loose when dry; non-sticky, non-plastic; strongly acid; clear, smooth boundary.
- BC1 14 to 23 inches, light yellowish brown (10YR 6/4, moist), sand; single grained; loose when moist, loose when dry; non-sticky, non-plastic; medium acid; gradual, wavy boundary.
- BC2 23 to 38 inches, light yellowish brown (10YR 6/4, moist), sand; single grained; loose when moist, loose when dry; non-sticky, non-plastic; medium acid; gradual, wavy boundary.
- C 38 to 48 inches, very pale brown (10YR 7/3, moist), sand; single grained; loose when moist, loose when dry; non-sticky, non-plastic; medium acid.

Analytical data in Table 10, Appendix I.

Forest Jack pine, M.A.I. = 26 cu.ft./ac./yr.

 $\frac{jP}{\text{Diameter}}$ Basal area 8.2 in. 60 ft. 130 yrs. 140 sq. ft.

Vegetation

Trees - Pinus banksiana

Picea mariana

Betula papyrifera

Tall shrubs - Rosa sp.

Shepherdia canadensis

Low shrubs - Arctostaphylos uva-ursi

Vaccinium angustifolium

V. vitis-idaea

Forbs - Anemone patens

Apocynum androsaemifolium

Aralia nudicaulis
Cornus canadensis
Geocaulon lividum
Linnaea borealis

Lycopodium spp.

Maianthemum canadense

Pyrola asarifolia

P. secunda

Capability Classification

Forestry: 55

Agriculture: 65

Wildlife: 55

Recreation: 60EQ

PLOT #9 Stereotriplet L Stereopair M Beach ridge on upper slope; gravelly calcareous sand.

Moisture regime: fresh (2), or well drained.

Soil Woodridge Complex

Land

The Woodridge Complex consists of rapidly to well drained Orthic Grey Luvisol, Orthic Eutric Brunisol

soils developed on stratified calcareous sand and gravel outwash and beach deposits.

- L-H 1½ to 0 inches, black (10YR 2/1, moist), forest litter; very strongly acid; abrupt, smooth boundary.
- Ae 0 to 1 inches, dark greyish brown (10YR 4/2, moist), loamy sand; single grained; loose when moist, loose when dry; non-sticky, non-plastic; medium acid; clear, smooth boundary.
- Bt 1 to 8 inches, very dark greyish brown (10YR 3/2, moist), loamy sand; weak, medium granular; very friable when moist, slightly hard when dry; non-sticky, non-plastic; neutral; mildly alkaline; gradual, wavy boundary.
- BC 8 to 12 inches, brown (10YR 5/3, moist), sand; single grained; loose when moist, loose when dry; non-sticky, non-plastic; mildly alkaline; clear, wavy boundary.
- C 12 to 20 inches, pale brown (10YR 6/3, moist), sand; single grained; loose when moist, loose when dry; non-sticky, non-plastic; moderately alkaline.

Analytical data in Table 11, Appendix I.

Forest Jack pine-black spruce, M.A.I. = 34 cu.ft./ac./yr.

	<u>jP</u>	<u>bS</u>
Diameter	8.2 in.	8.3 in.
Height	63 ft.	55 ft.
Age	92 yrs.	103 yrs.
Basal area	110 sq. ft.	30 sq. ft.

Vegetation

Trees - Pinus banksiana
Picea mariana

Tall shrubs - Alnus crispa

Rosa sp.

Low shrubs - Arctostaphylos uva-ursi

Ledum groenlandicum
Symphoricarpos sp.
Vaccinium vitis-idaea

Viburnum edule

Forbs - Cornus canadense
Lycopodium spp.
Petasites palmatus

Bryophytes & lichens

(On litter) Dicranum rugosum

D. drummondii

Pleurozium schreberi Hylocomium splendens

Brachythecium sp.

(In open area)

Ceratodon purpureus

(On rocks)

Hypnum reptile

Capability Classification

Forestry: 5™

Agriculture: 6 ₺

Wildlife: 55

Recreation: 60EQ

Stereopair 0. A19705:144-145, 1967. Scale ±1/16,000. T.64, R.24W. Detail of Stereopair N.

A13893:5-6, fall 1953. Scale ±1/68,000.

Stereopair N.

LEAK Landscape Unit (92 sq. mi.)

Stereotriplet

Weakly broken area of deep sandy till and shallow loamy till over dolomitic bedrock, with peat plains in depressions.

Lakes cover 8% of the area. Lakes of various size classes occupy the following proportion of the landscape unit:

Relatively small lakes: 3% Small lakes: 5%

The lakes are generally shallow and have somewhat irregular shorelines. The shore material is mainly bouldery till or, locally, organic matter. Most lakes have no open outlets and drain through bogs.

Land Unit ML-72L

Weakly broken area of knolls, slopes and flats of deep somewhat calcareous silty sand (70%), with plateaus of shallow highly calcareous loam over dolomitic bedrock (20%). Minor areas of deep mesic to fibric peat in depressions.

PLOT #10 Land Shallow fibric peat over dolomitic limestone.

Stereopair H&O Moisture regime: wet (7), or poorly drained.

Soil Flux Lake Series

The Flux Lake series consists of poorly to very poorly drained Lithic Fibric Mesisol soils developed on shallow (16 to 36 inches) forest peat or thin (\(\alpha \) 24 inches) sphagnum peat overlying forest peat.

Of - 0 to 16 inches, dark greyish brown (10YR 4/2, moist), non-woody, fibrous sphagnum peat; very strongly acid; fiber content approximately 100 percent.

Om - 16 to 23 inches, black (10YR 2/1, moist), moderately decomposed forest peat; neutral; fiber content approximately 54 percent.

R - 23+ inches, limestone bedrock.

Analytical data in Table 12, Appendix I.

Forest Black spruce, M.A.I. = 6 cu.ft./ac./yr.

bS

Diameter 4.7 in.

Height 25 ft.

Age 121 yrs.

Basal area 60 sq. ft.

Vegetation

Trees - Picea mariana

Larix laricina

Tall shrubs - Potentilla fruticosa

Salix bebbiana

Shepherdia canadensis

Low shrubs - Andromeda polifolia

Chamaedaphne calyculata

Ledum groenlandicum Vaccinium vitis-idaea

Forbs - Arctostaphylos rubra

Calla palustris

Equisetum sp.

Geocaulon lividum

2

Oxycoccos microcarpus

Bryophytes & lichens

(On litter) Meesia uliginosa

Pleurozium schreberi Tomenthypnum nitens

Aulacomnium palustre

Bryum pallens

Leptobryum pyriforme

Sphagnum nemoreum

Capability Classification

Forestry: 7 W

Agriculture: **7**^w**0**

Wildlife: 6 M 3 G
Recreation: 6 OER

LEAK Landscape Unit (92 sq. mi.)

Stereopair R

Weakly broken area of deep sandy till and shallow loamy till over dolomitic bedrock, with peat plains in depressions.

Lakes cover 8% of the area. Lakes of various size classes occupy the following proportion of the landscape unit:

Relatively small lakes:

Small lakes:

5%

3%

The lakes are generally shallow and have somewhat irregular shorelines. The shore material is mainly bouldery till or, locally, organic matter. Most lakes have no open outlets and drain through bogs.

Land Unit ML-74PL

Weakly broken area of deep somewhat calcareous silty sand slopes and flats (60%) and plateaus of shallow highly calcareous loam over dolomitic bedrock (20%). Deep mesic to fibric peat occurs in many depressions (20%).

Land Level dolomite plateau, bare to very shallow over PLOT #11 bedrock. Moisture regime: dry (0), or rapidly drained. Stereotriplet 2


> Rock Outcrop Soil

> > The Rock outcrop consists of 0 to 4 inches of rapidly drained surface deposits over limestone bedrock.

L-H - 0 to 2½ inches, very dark grey (10YR 3/1, moist), forest litter; slightly acid; abrupt, smooth boundary.

- 0+ inches, limestone bedrock. R

Analytical data in Table 13, Appendix I.



Stereotriplet Q. A19731:99-101, 1967. Scale ±1/16,000. T.65, R.23W. Detail of Stereopair O.

Forest Jack pine-black spruce, M.A.I. = 12 cu.ft./ac./yr.

bS jΡ tΑ 6.7 in. 8.0 in. 8.5 in. Diameter 40 ft. 43 ft. 45 ft. Height Age 140 yrs. 105 yrs. 110 yrs. Basal area 30 sq. ft. 30 sq. ft. 10 sq. ft.

Vegetation

Trees - Pinus banksiana
Picea glauca
Picea mariana
Populus tremuloides
Betula papyrifera

Tall shrubs - Juniperus communis

Lonicera sp.

Prunus virginiana

Rosa sp.

Shepherdia canadensis

Low shrubs - Arctostaphylos uva-ursi Juniperus horizontalis Viburnum edule

Forbs - Campanula rotundifolia
Galium trifidum
Linnaea borealis
Maianthemum canadense
Rubus pubescens
Viola spp.

Bryophytes & lichens

(On dolomite cliffs)

Myurella julacea

Encalypta rhabdocarpa

Ditrichum boreale

Hypnum fertile

Platygyrium repens

(On litter)

Dicrarella Sp


(On litter) Dicranella sp. Cladonia spp.

Capability Classification

Forestry: 6 R

Agriculture: 7 R

Wildlife: 6 Recreation: 6 Recreation: 6 OE



Scale ±1/68,000. T.64, R.21W. A13893:9-11, fall 1953. Stereotriplet R.

LEAK Landscape Unit (92 sq. mi.)

pest, neutral, Liber content approxim

Stereotriplet R

Weakly broken area of deep sandy till and shallow loamy till over dolomitic bedrock, with peat plains in depressions.

Lakes cover 8% of the area. Lakes of various size classes occupy the following proportion of the landscape unit:

Relatively small lakes:

3%

Small lakes:

5%

The lakes are generally shallow and have somewhat irregular shorelines. The shore material is mainly bouldery till or, locally, organic matter. Most lakes have no open outlets and drain through bogs.

Land Unit ML-72PL

Weakly broken area of somewhat calcareous silty sand slopes and flats (50%) and plateaus of shallow, highly calcareous loam over dolomitic bedrock (30%). Deep mesic to fibric peat occurs in many depressions (20%).

PLOT #12 Land Deep fibric peat on 4% slope.

Stereotriplet R


Stereopair S Soil Morin Lake Series

The Morin Lake series consists of poorly to very poorly drained Mesic Fibrisol soils developed on very deep (64 inches) sphagnum peat underlain by forest and/or fen peat. Moderately coarse sediments occur below 64 inches of the surface.

Of1 - 0 to 16 inches, black (10YR 2/1, moist), fibrous, sphagnum peat mixed with woody peat; slightly acid; fiber content approximately 72 percent.

- Of2 16 to 46 inches, black (10YR 2/1, moist), fibrous, sphagnum peat, neutral; fiber content approximately 81 percent.
- Om 46 to 71 inches, black (10YR 2/1, moist), moderately decomposed forest peat; neutral; fiber content approximately 60 percent.
- IIAhg 71+ inches, very dark greyish brown (10YR 3/2, moist), sand; single grained; mildly alkaline.

Analytical data in Table 14, Appendix I.





Stereopair S. A20044:161-162, 1967. Scale ±1/16,000. Reed Lake - T.64, R.24W.

Stereopair T. A13789:201-202. Scale ±1/16,000. T.63, R.26W.

APPENDIX I

Mechanical and Chemical Analysis of Soils

Table 1. Analysis of Fay Lake loamy sand

Ohemical Characteristics

	+H		09.97	2.30	1.14	0.74	92.0
tions	Na+		1.12	0.76	I. OI	0.77	0.52
able Ca 100 gms	K+		1.67	0.15	0.15	90.0	90.0
<pre>!xchangeable Cations m.e./100 gms.</pre>	Mg2+		5.15	0.93	0.34	0.23	0.43
H	Ga 2+		20.50	2.56	7.46	1.02	1.26
G.E.G.			81.90	09.9	3.98	2.72	2.36
C/M Ratio			62	37	10	1.1	ı
% Total N			0.89	0.03	70.0	0.02	ı
Org. C.			64.90	1.12	0.39	0.21	ı
Jolom, te			1	1	ı	1	1
69 150 Les		and supplied to the supplied of the supplied o	I	ı	1	ı	ι
Sold Funtar	3	And the bandon does the property of the star of a star or of	i	ı	i	ı	ı
Hd		-	75.35	100	5.70	5.70	6.35
Depth			2	\ c	100	10-01	24-29
Hor.			Et !	1 0	a d	RC	0

Particle Size Characteristics

Hor.	Depth	Stones	ro.		Sand 1	Fraction 2	on 3		Total	Silt	Clay	Textural
	inches	Size Size	2	V.C.S.	0	1 1	5.	V.F.S.	Sand	P6	P.S.	Class
Ex	2	ı	1	ı	1	ı	ı	1	ı	ı	ı	ı
1 0		1/1-1/2"	43.26	20.54	14.62	13.76	15.31	14.36	78.59	17.34	4.07	L.S.
	C - 7	1/1-1/3"	17.77	2002	13.72	16.37	22.13	18,21	75.47	19.72	4.81	I.S.
	10101	1/1-1/2	37.53	76.8	16.11	17.4.7	21.62	15.56	79.67	17.09	3.24	H.S.
	57-20	1/1-3"	77	7.80	15.18	18.77	20.38	17.03	79.10	17.66	3.24	.,

Table 2. Analysis of Rock Outcrop (Precambrian)

Hor.	Depth inches	hd	Cond. mmhos/ cm	CaCO3 Equiv.	Cal-	Jolo- mite	Org.	Total N	C/N Ratio	G.E.C.	Exchangeable Cations m.e./100 gms.	geable /100 g	Cation Zms. Na+	H+
L-H	5	5.20	0.55	ı	ı	i	22.83	06.0	25	96.00	20.95 7.25 0.83 0.35 19.15	0.83	0.35	19.15

Table 3.
Analysis of Athapap sandy loam

Chemical Characteristics

Cond.	en c				0.16	
	H+	•	43.01	9.55	2.46	0.79
ations 3.	Na.		0.33	0.28	0.21	0.22
100 gm	K+		1.92	0.13	0.10	0.11
Exchangeable Cations m.e./100 gms.	• Mg ² +		6.70	1.45	0.65	0.25
1	ca2+		20.25	1.80	1.25	1.70
C.E.C.			80.70	15.20	6.10	4.95
C/N Ratio			43	77.	19	
% Total N			1.32	0.07	0.03	1
% Org. G.			56.3	1.66	0.58	ı
% Dolomite			1	1	1	1
% V. Calcite D			ı	ı	1	1
Saco Equiv.			1	1	1	1
Hď		il.	7.35	7.70	08.7	5.25
Depth			73.0	7 1	9 1 7	6 -12
Hor.			H	i E	BGo	, p

Particle Size Characteristics

lor.	Depth	Stones	Ø		Sand	Sand Fraction %	% uo		Total	Silt	Clay	Textura
	inches	Dominant Size	BS	V.C.S.	c.s.	M.S.	F.S.	V.F.S.	Sand	se	se	Class
ш	13.	,	ı	ı	ı	1	1	1	1	1	1	
	7 -0	1/3-1/4"	27.27	10.45	. 12,48	10.98	10.33	8.86	53.09	35.76	11.15	S.L.
Co	9 1 7	1/2-3/4"	18.37	6.10	10.82	11.39	13.54	13.07	54.92	36.76	8,32	
0 5d	6 -14	1/4-1/3"	21.72	8,00	11.84	11.23	13.63	13.44	58.14	33.53	8.33	

Table 4.

Analysis of Herb Lake clay

Chemical Characteristics

Cond.	uo	0.27	0.18	0.15	0.20
	#H	18.97	6.47	3.45	4.28
ıtions	Na+	0.71	0.31	0.23	1.89
able Ga 100 gms	K+	1.09	0.22	0.31	0.40
Exchangeable Cations m.e./100 gms.	Mg2+	6.20	3.08	5.75	9.55
H	Ca2+	25.00	3.88	9.75	14.00
C.E.C.		\$2.90	16.90	21,00	27, 80
c/N Ratio		89	77	17	17
% Total N		0.93	90.0	0.07	0.05
org. C.		63.45	1.46	1.01	0.85
% Dolomite		ı	ı	ı	1
% Calcite		1	1	ı	1
% CaCO ₃ Equiv.	e	ı	1	.1	1
Hď		7.40	4.55	5.15	5.30
Depth inches		3-0	0 <mark>-</mark> 3	3-5	5-10
Hor.		I-H	Ae	AB	Btg

Particle Size Characteristics

Hor.	Depth	Stones	ω.		Sand	Sand Fraction %	% uo		Total	Silt	Clay	Textural
-	inches	Dominant Size	مح	V.C.S.	c.s.	M.S.	F.S.	V.F.S.	Sand	<i>b</i> e	pe .	Class
E	9	ı	ı	/ ₁ 1	1	1	1	1	ı	J	1	1
Ae	9	1/3-1/2"	14.55	2.90	6.67	6.31	6.62	99.9	29.15	37.63	33.22	C.L.
AB	7 5	₩	6.88	2.34	3.95	4.25	14.72	3.93	19.19	31.08	49.73	ن
Btg	5-10	1/3-1/2"	7.42	1.58	3.43	4.05	4.47	07.7	17.93	24.22	57.84	·0

Table 5. Analysis of Batty Lake Series (Gryic Mesic Fibrisol)

Hor.	Depth Inches	% Un- rubbed	Pyro-	% Ash	% % Ash Org.	of Total	c/M Ratio	Hq ur	G.H.G.	Ä	Exchangeable Cations m.e./100 gms.	hle Ca 00 gms	tions	
		ri ber			ی	Z.		KCL		0s2+	Mg2+	K+	Mg ²⁺ K+ Ma+ H+	±
J 0	0- 37	72.9	0.092	7.83	7.83 56.1 1.11	1.11	50.5 4.65	4.55	6.701	60.80	20.60	69.0	20.60 0.69 0.51 41.5	41.5
Omz	37-170	6.95	0.076	ı	50.0	50.0 1.65	30.3 5.30	5.30	60.1	73.32	18.76 0.25 0.61	0.25	0.61	27.8
Om	1.70+	ı	ı	I	٠. 1	ı	ı	ı	1	1	ı	ı	ı	ı

Table 6. Analysis of Egg Lake sandy loam

Chemical Characteristics

	##		0	0	1.2	ŧ	ı	1	
ations 3.	Na+	(7.0	&. ○	2.0	ı	ı	ı	
able Ca	K+		0.0	0.1	0.3	ı	1	ı	
Exchangeable Cations m.e./100 gms.	Mg ² +	1	T2.0	1.4	3.0	ı	ı	ı	
	Ca ²⁺		49.4	3.6	7.1	ı	ı	ı	
C.E.C.			3.79	4.5	12,8	ı	ı	1	
C/N Batio			77	14	17	ı	ı	ı	
8, Total N	7	1	1.25	0.03	0.03	1	ı	ı	
Oro. G.			29.58	0.43	0.51	ı	ı	ı	
% 6, FOUT - FOUT			1	ı	1	47.9	7.87	00.9	
8, 0, 10, 10, 10, 10, 10, 10, 10, 10, 10,	7075		ı	ı	ı	6.20	12.18	6.52	
H S ODED	Jan 3 Harry		ı	1	1	13.19	20.72	13.04	
Hď		1			6.20				
Depth	apriorita		19-0	8 10	8 -14	14-19	19 -34	34+	
Hor.			LH	Ae	3 t	BC	Cca	O	

Particle Size Characteristics

Hor.		Stones	,		Sand	Sand Fraction %	% uo		Total	Silt	Clay	Textural
	inches	Dominant Size	P.J.	V.C.S.	ນ ໝ	M.S.	F4 S2	V.F.S.	Sand	₽ć.	₽6	Class
3	13-0		ı	1	1	. 1	1	1		1	ı	ı
Ae	8 1 0	ı	1	6.15	12.45	13.75	16.43	17.51	66.29	25.35	8.35	S.L.
34	8 -14	1	1	5.49	9.54	11.83	14.03	13.21	54.10	27.42	18.48	S.L.
33	14-19	1	ı	4.27	10.03	11.75	15.53	13.53	55.11	31.37	13.53	S.L.
Jea.	19 -34		ı	13.21	13.05	11.18	12.21	11.83	61.47	26.10	12.42	S.L.
-	34+	1	1	5.98	10.78	11.70	13.10	13.97	55.54	32.45	13.02	S.L.

Table 7. Analysis of Batty Lake Sories (Gryic Mesic Fibrisol)

1	Ĭ	. 1					bbb
	% % 11- Dolo-	mite	ı	1	0.0		13.4
	Cal-	cite	1	1	1.81 0.0		
-	Gaco ₃	100	1	1	1.81		
		+	1	1	1		
	ions	Na+	0.23	0.30	0.37		
-	le Cat	K+	0.71	0.20	0.25		
	"xchangeable Cations m.e./100 gms.	Mg ²⁺	16.36	15.15	23.63	a y	c 1 a y
	TXC.	ca ² +	39.9 6.12 180.2 153.11 16.36 0.71 0.23	151.29 15.15 0.20 0.30	157.35 23.63 0.25 0.37	e clay	
	C.E.C.		180.2	23.3 6.88 210.0	28.5 7.20 194.8	lacustrine	en lacustrine
	Hq.	TON I	6.12	6.88	7.20	c u s	a c u
	c/N Ratio		39.9	23.3	28.5		n 1
	% % C/N pH Org. Total Ratio in	2	1.12	33.8 1.88	1.83	ozen	12
	org.	5	7.47	43.8	52.1 1.83	froz	unfro
	Ash		15.59 44.7 1.12	ı	ı		
	Pyro-	phos.	0.592	0.275	0.201		
-	% Un- % rubbed Pyro-	Fiber	0- 7 81.80	7- 34 81.0	42.1		
		15	7	37	34- 96	63	191
	Depth inches		٩	7-	34-	96-163	163-167
	Hor.		0£1	0£2	Omz	IICgz	IICg

Table 8. Analysis of Wabowden clay

Chemical Characteristics

Con	CB		7.0	0	0.1	0.1	0	7.0	0
	₩		34.85	5.20	3.56	2,62	1	ı	ı
ations s.	Na+		1.54	0.97	1.10	1.28	ı	ı	ı
Exchangeable Cations m.e./100 gms.	K+		2.95	0.00	1.13	0.97	ı	ı	ı
Exchang m.e.	Mg^{2}		10.85	8.80	12,60	11.72	ı	ı	1
	Ca2+		22.63	10.08	17,40	19.44	ı	ı	ı
C.E.C.			49.92	25.16	29.86	32.39	30.44	25.75	22.12
C/N Ratio			26	77	12	10	10	1	1
% Total N		3	0.95	0.11	0.08	90.0	0.05	1	ı
% Org. C.	2		52.75	1.53	0.92	0.61	0.52	ı	1
% te Dolomite	1		ı	ı	ı		0.0	3.33	5.81
% Calcite			ı	ı	ı	ı	1.07	1.66	15.19
% CaCO ₃ Equiv.			1	1	1	ı	1.07	5.28	21.50
Hd		8					7.10		
Depth inches			5 · 0	0 - 43	42- 92	92-183	183-223	223-30	30 -39
Hor.		:	L	Ae	AB	Bt	BC	5	22

Particle Size Characteristics

Hor.	Depth	Stones			San	d Fract	ion %		Total	Silt Clay	Clay	Textural
88	inches	Dominant Size	pe	1- 1	c.s.	M.S.	દ્ય જ	7.C.S. C.S. M.S. F.S. V.F.S.	Sand	6 9	<i>6</i> 2	Class
LH	2 - 0	l	1	I	5.1		; I	1	-	•		
Ae	0 - 43	I	ı	1	1	1	ı	ı	3.84	26.97	69.19	5
AB	43- 93	I	i	ı	1	ı	I	ı	ı	1	ı	ల
Bt	95-183	l	ı	ı	ı	ı	ı	1	ı	ı	ı	ల
BC	183-223	I	1	ı	1	ı	ı	1	1.54	19.25	79.21	ပံ
15	223-30		1		1	1	ı	1	2,48	27.66	98.69	ల
8	30 - 39	1	ı	. 1	ı	ı	ı	1	2.43	33.69	63.88	Ü

Table 9.

Analysis of Dering fine sandy loam

Chemical Characteristics

Cond.	e c		0.24	0.32	0.33	0.39
	H+		52.90	2.80	1.15	ı
tions	Na+		1.46	1.57	1.77	1
able Ca 100 gms	K+		4.42	1.47	0.29	ı
Exchangeable Cations m.e./100 gms.	Mg ² +		18.27	64.45	22.25	ı
	Ca2+		28.90	56.80	52,00	ı
C.E.C.			112,05	120.80	70.40	11.44
c/N Ratio	1		ı	147	77	ı
% C. Total N R			0.56	1.19	0.55	1
Org. C.			58.79	56.20	13.45	ı
% Dolomite			1	1	1	22.64
% Calcite			1	ı	1	15.18
% CaCO ₂ Equiv.			1	ı	1	39.77
		100	4.30	6.90	6.80	7.50
Depth pH			83-43	43-0	7-0	4-11
Hor.			L-F	Н	Ahg	် ရှိ

Particle Size Characteristics

lor.		Stones	ro		Sand	Sand Fraction %	% uo		Total	Silt Clay	Clay	Textural
	inches	Dominant Size	88	IP	. C. S. C. S.	M.S. F.S.	بر دي دي	V.F.S.	Sand	90	96	Class
E4	83-43	1	ı	ı	1	1	ı	1	1	1	1	1
	0-47	1	ı	ı	1	1	1	1	1	ı	1	ı
ha	7-0	1	ı	3,80	5.53	99.6	25.11	10,99	55.09	30.44	14.46	F.S.L.
Cas	4-11	ı	ı	2.68	3.33	4.51	7.33	6.91	24.74	51.74	23.52	Si.L.

Table 10 Analysis of Clarke sand

Chemical Characteristics

Hor.	Depth inches	Hď	CaCO3 Equiv.	Calc	% Dolomite	Org. C.	% Total N	C/N Ratio	G.H.G.		Exchange m.e./	Exchangeable Cations m.e./100 gms.	ations		Cond. mmhos/
										Ca2+	Mg ² +	K+	Na+	H	cm
		15										1		7	
L-H	2-0	2.00	1	ı	ı	27.71	0.82	34	65.80	37.93	23.28	1.60	1.33	0.85	99.0
Ae	9 -0	5.20	1	1	ı	0.53	0.02	27	2.08	79.0	0.72	90.0	0,40	0.30	0.13
Bm	41−9	5.50	1	1	L	0.19	0.01	19	2.22	0.94	0.03	0.05	0.75	0.70	0.10
BC1	14-23	6.05	1	ı	1	0.18	0.01	18	1.80	1.40	0.24	0.08	0.85	0.24	0.07
BC2	23-38	6.20	1	ı	ı	0.10	0.01	10	1.60	1.10	0.38	90.0	0.57	00.00	0.11
Ö	38-48	7.20	6.93	%*0	5.50	1	ı	1	1.02	1	ı	ı	1	ı	0.14

Particle Size Characteristics

Hor.	Depth	Stones	ω		Sand	Sand Fraction %	% uo		Total	Silt	Clay	Textural
28	inches	Dominant Size	<i>6</i> 2	V.C.S.	o	M.S.	F. S.	V.F.S.	Sand	6 9	96	Class
L-H	2-0	ı	ı	ı	7	1	1	,	. 1		1	1
Ae	9-0	-	1	1	4.46	14.37	50.80	25.46	60.56	4.10	0.81	ß
Bm	6-14	1	1	ı	0.79	11.47	56.95	26.34	95.54	4.37	0.09	ß
BC1	14-23	C	ı	I	69.0	11.51	26.44	27.20	78.56	2.00	2.16	S
BC2	23-38	. 1	i	1	1.71	35.84	47.81	12.82	98.18	87.0	1.34	co
O	38-48	1	1	1	1.76	28.31	52.21	14.32	09.80	3.02	0.38	ß

Table 11.
Analysis of Woodridge gravelly sand

Chemical Characteristics

Cond.	cm	0.31	0.38	0.58	0.33	0.32
	+H	33.10	5.25	1	ı	ı
ations 18.	Na+	1.63	0.0	ı	ı	I
Exchangeable Cations m.e./100 gms.	+ X	1.86	0.19	ı	ı	ı
Exchang m.e.	Mg ² +	20.15	3.52	1	ı	ı
	Ca2+	25.40	4.43	ı	ı	ı
C.E.C.		87.40	14.72	18.60	4.02	3.00
C/N Ratio		87	ı	18	7	ı
% Total N		1.23	1	0.19	0.04	ı
org. G.		58.54	4.50	3.50	0.29	
% Dolomite		ı	ı	41.71	47.95	34.77
% Calcite		1	ı	0.0	7.23	5.35
CaCO ₃ Equiv.		1	ı	45.28	59.29	43.09
Hd		4.85	7.80	7.40	7.65	7.90
Depth		13-0	0 - 1	1 8	8 -12	12 -20
Hor.		L-H	Ae	Bt	BC	O

Particle Size Characteristics

Hor.	Depth	Stone	ω		Sand	Sand Fraction %	% uo		Total	Silt	Clay	Textura]
	inches	Dominant Size	83	V.C.S.	S.S.	M.S.	સ જ.	V.F.S.	Sand	66	82	Class
Ę	13-0	ı	ı	ı	ı	. 1	ı	ı	ı	ı	1	1
Ae	0 - 1	1/3"	3.90	7.85	28,66	18.74	14.13	8,62	77.99	15.12	68.9	L.S.
Bt	1 - 8	1/2-1"	17.77	10.75	54.49	69.63	6.03	24.17	75.07	20.93	7.00	L.S.
BC	8 -12	1/4-1/3"	37.46	12.68	47.45	20.68	2.77	6.41	86.68	9.73	0.29	'n
O	12 -20	1/2-3/4"	55.74	12.70	38.51	27.47	10,66	5.06	94.41	5.38	0.21	so.

Table 12.
Analysis of Flux Lake Series (Lithic Fibric Mesisol)

	+ H	1.15	0.30	
tions	Na+	0.03	0.04	
ble Ca	*	0.08	0.05	
Exchangeable Cations m.e./100 gms.	Ca ²⁺ Mg ²⁺ K+ Na+	37.08	78.60	
Exc	ca ²⁺	90.96 37.08 0.08 0.03 1.15	135.27 48.60 0.05 0.04 0.30	
C.E.C.		4.85 127.5	7.10 204.5	٥ د د
Hd in	TOU	4.85	7.10	bedrock
C/N Ratio		58	32	
Total		99.0	1.12	limestone
8 4 0 c		7.25 38.0 0.66	36.0	₽ H
% Ash		7.25	13.25 36.0 1.12	H
Pyro-		0.087	0.458	
Depth % Un- inches rubbed	10011	100	54.5	
		0-16	16-23	23+
Hor.		JO	Om	ಚ

Table 13.
Analysis of Rock Outcrop (limestone)

Chemical Characteristics

Cond.	u	32	
Col	CH	0	
	†H	00°α	
Exchangeable Cations m.e./100 gms.	Na+	1.24	
eable (K+	0.90	
Exchang m.e.	11g2+	14.50	
	Ca2+ 11g2+ K+ Na+	33.20	
G.E.G.		38 61.01 33.20 14.50 0.90 1.24 2.00 0.35	o Y
C/N Batio		38	9 d r
% 6/M C.E.C.		0.55	mestone Bedrock
<i>b</i> 9		20.69 0.55	α t
		20	E E
Salcite Dolomite		1	r
8 Ca	4	1	
CACO Hourty		I	
		6.10	
Depth pH		т-н 0-23 6.10	
Hor.		1 -H	æ

Table 14.
Analysis of Morin Lake Series (Mesic Fibrisol)

Hor.	Depth inches		Mechanical Analysis	. Analys	\$1.	% Un-	P 20	16 2	25 24	5€ + €	C/N		C.E.C.	DXC	Exchangeable Cations	ble Ca	tions	1
			Text. % % % %	Silt	% Clay	Fiber	phos.	HON I	0	N		KCl	Ф	Ca2+	Ca2+ Mg2+ K+ Na+	K+ K+	Na+	压
)£1	0-16	1	1	1	1	71.6	0.231	77.54	06.0 36.5 75.11	06.0	5	7 7	8 /01 71 9	ر /۶۲	000	6	3	1 6
260	7/-71						1 (1 0) (•) ·		174.5 44.07 0.03 0.04 0.1	44.03	0.0	40.0	
717	1040	ı	ı	ı	ı	×0.08	0.113	88.6	36.8	0.73	20	06.9	190.50	150.30 46.84 0.02 0.05 0.5	78.97	0.02	0.05	7.0
Jm	TL-9 [†] 7	ı	ı	1	1	61.3	0.885	16.17 34.4	34.4	1.04	33	7.30	183.3	142,30	45.48 0.03 0.06	0.03	90.0	1
IAhg	477	S.L.	44.32	44.32 47.78 7.90	7.90	ı	ı	ı	10.8 0.64	79.0	17	7.80	21.80	7.80 21.80 120.24 9.56 0.02 0.02	9.56	0.02	0.02	ı